

Determination of $2\beta_s$ in $B_s^0 \rightarrow J/\psi K^+ K^-$ Decays in the Presence of a $K^+ K^-$ S-Wave Contribution

Yuehong Xie,^a Peter Clarke,^b Greig Cowan^c and Franz Muheim^d

*School of Physics and Astronomy, University of Edinburgh,
Mayfield Road, Edinburgh, EH9 3JZ, U.K.*

Abstract

We present the complete differential decay rates for the process $B_s^0 \rightarrow J/\psi K^+ K^-$ including S-wave and P-wave angular momentum states for the $K^+ K^-$ meson pair. We examine the effect of an S-wave component on the determination of the CP violating phase $2\beta_s$. Data from the B-factories indicate that an S-wave component of about 10% may be expected in the $\phi(1020)$ resonance region. We find that if this contribution is ignored in the analysis it could cause a bias in the measured value of $2\beta_s$ towards zero of the order of 10%. When including the $K^+ K^-$ S-wave component we observe an increase in the statistical error on $2\beta_s$ by less than 15%. We also point out the possibility of measuring the sign of $\cos 2\beta_s$ by using the interference between the $K^+ K^-$ S-wave and P-wave amplitudes to resolve the strong phase ambiguity. We conclude that the S-wave component can be properly taken into account in the analysis.

^aYuehong.Xie@cern.ch

^bpeter.clarke@ed.ac.uk

^cg.cowan@ed.ac.uk

^df.muheim@ed.ac.uk

Contents

1	Introduction	1
2	Time-dependent angular distributions in the decay $B_s^0 \rightarrow J/\psi K^+ K^-$ including S-wave contributions	1
3	Measuring $2\beta_s$ in the presence of a $K^+ K^-$ S-wave	4
4	Measuring $\cos 2\beta_s$	7
5	Conclusions	8

1 Introduction

The decay $B_s^0 \rightarrow J/\psi \phi$ is a golden channel for the measurement of the B_s^0 mixing phase $-2\beta_s$ which is a very sensitive probe of new physics. It has been extensively studied [1, 2, 3, 4, 5, 6, 7, 8]. In the decay $B_s^0 \rightarrow J/\psi \phi$, followed by a two-body decay $\phi(1020) \rightarrow K^+ K^-$, the $K^+ K^-$ meson pair is in an orbital P-wave amplitude. However, in the vicinity of the $\phi(1020)$ mass, the $K^+ K^-$ system can have contributions from other partial waves. The same comment holds for the $K^+ K^-$ system in the decay channels $B^0 \rightarrow K^+ K^- K_S^0$ and $D^0 \rightarrow K^+ K^- \pi^0$. The BaBar experiment showed that in these decays the S-wave and P-wave contributions dominate in the mass range above threshold up to $1.1 \text{ GeV}/c^2$ [9, 10]. In both cases there is a dominant resonant $\phi(1020)$ contribution. In addition an S-wave $f_0(980)$ and a non-resonant contribution are found to be necessary to describe the data. These results motivated us to investigate the effects of a possible S-wave contribution to $B_s^0 \rightarrow J/\psi K^+ K^-$ in the $\phi(1020)$ mass region.

In the decay $B_s^0 \rightarrow J/\psi K^+ K^-$ the $K^+ K^-$ system can only arise from a $s\bar{s}$ quark pair while in $B^0 \rightarrow K^+ K^- K_S^0$ and $D^0 \rightarrow K^+ K^- \pi^0$ it can have contributions from both $s\bar{s}$ and $d\bar{d}$. This makes it difficult to give a quantitative estimate for the S-wave component. In reference [11] the S-wave $K^+ K^-$ contribution under the $\phi(1020)$ peak is estimated to be $5 - 10\%$ for decay modes in which the $K^+ K^-$ arises from an $s\bar{s}$ quark pair. In this study we consider an S-wave of similar magnitude and assess its impact on the determination of the weak mixing phase $-2\beta_s$.

2 Time-dependent angular distributions in the decay $B_s^0 \rightarrow J/\psi K^+ K^-$ including S-wave contributions

We consider P- and S-wave amplitudes in the decay $B_s^0 \rightarrow J/\psi K^+ K^-$ where the invariant mass of the $K^+ K^-$ meson pair is in the $\phi(1020)$ mass region and the J/ψ meson decays into a

$\mu^+\mu^-$ pair. The S-wave contribution can be non-resonant or due to the $f_0(980)$ resonance¹. We denote decay amplitudes for the $B_s^0 \rightarrow J/\psi K^+ K^-$ by $\mathbf{A} = (A_0, A_{||}, A_{\perp}, A_S)$. Here A_0 , $A_{||}$ and $A_{\perp}$ are the three P-wave amplitudes consistent with the $K^+ K^-$ system decaying via the $\phi(1020)$ resonance. A_S is the amplitude for a possible S-wave contribution in the $K^+ K^-$ system. The amplitudes for the conjugate decay $\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$ are denoted by $\bar{\mathbf{A}} = (\bar{A}_0, \bar{A}_{||}, \bar{A}_{\perp}, \bar{A}_S)$, which, in the absence of direct CP violation, are related to $\mathbf{A}$ by $A_0 = \bar{A}_0$, $A_{||} = \bar{A}_{||}$, $A_{\perp} = -\bar{A}_{\perp}$ and $A_S = -\bar{A}_S$. Note that A_0 and $A_{||}$ are CP-even whereas $A_{\perp}$ and A_S are CP-odd. The amplitudes $(A_0, A_{||}, A_{\perp})$ and the amplitude A_S may have different dependences on the mass $m_{K^+ K^-}$ of the $K^+ K^-$ system. However, in sufficiently small bins of $m_{K^+ K^-}$, such as the narrow mass region around the $\phi(1020)$ resonance, the dependences of the amplitudes on $m_{K^+ K^-}$ can be neglected.

We define the total P-wave strength, $A_P^2 \equiv |A_0|^2 + |A_{||}|^2 + |A_{\perp}|^2$, the longitudinal and perpendicular polarisation fractions relative to the P-wave strength $R_{||} \equiv |A_{||}|^2/A_P^2$, and $R_{\perp} \equiv |A_{\perp}|^2/A_P^2$, and the S-wave fraction, $R_S \equiv |A_S|^2/(A_P^2 + |A_S|^2)$. The phases of these decay amplitudes are defined by $A_j = |A_j|e^{i\delta_j}$, where $j = 0, ||, \perp, S$. As only the relative strong phase differences can be measured we adopt the convention $\delta_0 = 0$.

An angular analysis is required to disentangle the different CP eigenstates on a statistical basis. The angular observables are denoted as the helicity angles $\Omega = (\theta_l, \theta_K, \varphi)$. Here θ_l is the angle between the μ^+ momentum and the direction opposite to the B_s^0 momentum in the J/ψ rest frame; θ_K is the angle between the K^+ momentum and the direction opposite to the B_s^0 momentum in the rest frame of the $K^+ K^-$ system; φ is the angle between the decay planes of the $J/\psi \rightarrow \mu^+ \mu^-$ and the $K^+ K^-$ pair, when going from the positive kaon to the positive lepton with a rotation around the opposite direction of the B_s^0 momentum in the J/ψ rest frame.

The differential decay rate for a B_s^0 meson produced at time $t = 0$ decaying as $B_s^0 \rightarrow J/\psi K^+ K^-$ at proper time t is given by

$$\frac{d^4\Gamma(B_s^0 \rightarrow J/\psi K^+ K^-)}{dt d\cos\theta d\cos\psi d\varphi} \propto \sum_{k=1}^{10} h_k(t) f_k(\Omega), \quad (2.1)$$

whereas the differential decay rate for an initial $\bar{B}_s^0$ meson is given by

$$\frac{d^4\Gamma(\bar{B}_s^0 \rightarrow J/\psi K^+ K^-)}{dt d\cos\theta d\cos\psi d\varphi} \propto \sum_{k=1}^{10} \bar{h}_k(t) f_k(\Omega). \quad (2.2)$$

Each of the $h_k(t)$, $\bar{h}_k(t)$ and $f_k(\Omega)$ for $k = 1 - 10$ are defined in Table 1. In total there are four amplitude-squared terms for the three polarisations of the P-waves and the S-wave component plus six interference terms.

The time-dependence of the ten functions $h_k(t)$ for an initial B_s^0 meson state can be written as:

$$|A_0(t)|^2 = |A_0|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) + \sin\Phi \sin(\Delta m_s t) \right], \quad (2.3)$$

$$|A_{||}(t)|^2 = |A_{||}|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos\Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) + \sin\Phi \sin(\Delta m_s t) \right], \quad (2.4)$$

¹The mass dependence of the $f_0(980)$ is distorted as the central value of the resonance is below threshold.

k	$h_k(t)$	$\bar{h}_k(t)$	$f_k(\theta_l, \theta_K, \varphi)$
1	$ A_0(t) ^2$	$ \bar{A}_0(t) ^2$	$4 \sin^2 \theta_l \cos^2 \theta_K$
2	$ A_{\parallel}(t) ^2$	$ \bar{A}_{\parallel}(t) ^2$	$(1 + \cos^2 \theta_l) \sin^2 \theta_K - \sin^2 \theta_l \sin^2 \theta_K \cos 2\varphi$
3	$ A_{\perp}(t) ^2$	$ \bar{A}_{\perp}(t) ^2$	$(1 + \cos^2 \theta_l) \sin^2 \theta_K + \sin^2 \theta_l \sin^2 \theta_K \cos 2\varphi$
4	$\Im\{A_{\parallel}^*(t)A_{\perp}(t)\}$	$\Im\{\bar{A}_{\parallel}^*(t)\bar{A}_{\perp}(t)\}$	$2 \sin^2 \theta_l \sin^2 \theta_K \sin 2\varphi$
5	$\Re\{A_0^*(t)A_{\parallel}(t)\}$	$\Re\{\bar{A}_0^*(t)\bar{A}_{\parallel}(t)\}$	$-\sqrt{2} \sin 2\theta_l \sin 2\theta_K \cos \varphi$
6	$\Im\{A_0^*(t)A_{\perp}(t)\}$	$\Im\{\bar{A}_0^*(t)\bar{A}_{\perp}(t)\}$	$\sqrt{2} \sin 2\theta_l \sin 2\theta_K \sin \varphi$
7	$ A_S(t) ^2$	$ \bar{A}_S(t) ^2$	$\frac{4}{3} \sin^2 \theta_l$
8	$\Re\{A_S^*(t)A_{\parallel}(t)\}$	$\Re\{\bar{A}_S^*(t)\bar{A}_{\parallel}(t)\}$	$-\frac{2}{3}\sqrt{6} \sin 2\theta_l \sin \theta_K \cos \varphi$
9	$\Im\{A_S^*(t)A_{\perp}(t)\}$	$\Im\{\bar{A}_S^*(t)\bar{A}_{\perp}(t)\}$	$\frac{2}{3}\sqrt{6} \sin 2\theta_l \sin \theta_K \sin \varphi$
10	$\Re\{A_S^*(t)A_0(t)\}$	$\Re\{\bar{A}_S^*(t)\bar{A}_0(t)\}$	$\frac{8}{3}\sqrt{3} \sin^2 \theta_l \cos \theta_K$

Table 1: Definition of the functions $h_k(t)$, $\bar{h}_k(t)$ and $f_k(\theta_l, \theta_K, \varphi)$ of Eq. 2.1 and 2.2.

$$|A_{\perp}(t)|^2 = |A_{\perp}|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + \cos \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin \Phi \sin(\Delta m_s t) \right], \quad (2.5)$$

$$\begin{aligned} \Im\{A_{\parallel}^*(t)A_{\perp}(t)\} &= |A_{\parallel}||A_{\perp}| e^{-\Gamma_s t} \left[-\cos(\delta_{\perp} - \delta_{\parallel}) \sin \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. + \sin(\delta_{\perp} - \delta_{\parallel}) \cos(\Delta m_s t) - \cos(\delta_{\perp} - \delta_{\parallel}) \cos \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.6)$$

$$\begin{aligned} \Re\{A_0^*(t)A_{\parallel}(t)\} &= |A_0||A_{\parallel}| e^{-\Gamma_s t} \cos(\delta_{\parallel} - \delta_0) \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) - \cos \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. + \sin \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.7)$$

$$\begin{aligned} \Im\{A_0^*(t)A_{\perp}(t)\} &= |A_0||A_{\perp}| e^{-\Gamma_s t} \left[-\cos(\delta_{\perp} - \delta_0) \sin \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. + \sin(\delta_{\perp} - \delta_0) \cos(\Delta m_s t) - \cos(\delta_{\perp} - \delta_0) \cos \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.8)$$

$$|A_S(t)|^2 = |A_S|^2 e^{-\Gamma_s t} \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + \cos \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) - \sin \Phi \sin(\Delta m_s t) \right], \quad (2.9)$$

$$\begin{aligned} \Re\{A_S^*(t)A_{\parallel}(t)\} &= |A_S||A_{\parallel}| e^{-\Gamma_s t} \left[-\sin(\delta_{\parallel} - \delta_S) \sin \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. + \cos(\delta_{\parallel} - \delta_S) \cos(\Delta m_s t) - \sin(\delta_{\parallel} - \delta_S) \cos \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.10)$$

$$\begin{aligned} \Im\{A_S^*(t)A_{\perp}(t)\} &= |A_S||A_{\perp}| e^{-\Gamma_s t} \sin(\delta_{\perp} - \delta_S) \left[\cosh\left(\frac{\Delta\Gamma_s t}{2}\right) + \cos \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. - \sin \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.11)$$

$$\begin{aligned} \Re\{A_S^*(t)A_0(t)\} &= |A_S||A_0| e^{-\Gamma_s t} \left[-\sin(\delta_0 - \delta_S) \sin \Phi \sinh\left(\frac{\Delta\Gamma_s t}{2}\right) \right. \\ &\quad \left. + \cos(\delta_0 - \delta_S) \cos(\Delta m_s t) - \sin(\delta_0 - \delta_S) \cos \Phi \sin(\Delta m_s t) \right], \end{aligned} \quad (2.12)$$

where $\Phi = -2\beta_s$, Δm_s , $\Delta\Gamma_s$ and Γ_s denote the weak mixing phase, mass difference, decay width difference and average decay width of the B_s^0 - $\bar{B}_s^0$ system, respectively. Here we have assumed that each of the decay amplitudes in $\mathbf{A}$ is dominated by a single weak phase, therefore a common effective $2\beta_s$ can be used for all CP eigenstates. The time evolution functions $\bar{h}_k(t)$ for an initial $\bar{B}_s^0$ meson can be obtained by reversing the sign of each term proportional to $\sin(\Delta m_s t)$ or $\cos(\Delta m_s t)$ in $h_k(t)$.

3 Measuring $2\beta_s$ in the presence of a K^+K^- S-wave

In this section we investigate how the measurement of $2\beta_s$ is affected by the presence of a possible K^+K^- S-wave contribution. We use Monte Carlo simulated toy data based on the differential decay rate expressions of Section 2. We generate signal decays only and ignore backgrounds underneath the B_s^0 mass peak as well as all detector effects. The inclusion of these effects does not alter the qualitative results of this study.

We assume a tagging efficiency $\epsilon_{tag} = 56\%$ and a wrong tag probability $\omega_{tag} = 33\%$, which correspond approximately to the expected flavour tagging performance for this channel at the LHCb experiment [12]. In Table 2 we summarize the values of the physical parameters used to generate the toy data sets.

We generate 500 data sets for different scenarios where we vary the values of the S-wave fraction R_S and its phase δ_S and the weak phase $-2\beta_s$. Each data set contains 30000 signal events corresponding to approximately one quarter of a nominal LHCb year of 2 fb^{-1} .

	Δm_s	Γ_s	$\Delta\Gamma_s$	δ_0	$\delta_{ }$	$\delta_{\perp}$	$R_{ }$	$R_{\perp}$	R_S	δ_S	$2\beta_s$
Input	17.8 ps^{-1}	0.68 ps^{-1}	0.05 ps^{-1}	0	-2.93	2.91	0.207	0.233	vary	vary	vary
Fit	fix	fix	fix	fix	float	float	float	float	float*	float	float

* R_S is fixed to 0 when the S-wave component is neglected.

Table 2: Values of the physical parameters used in the generation of signal decays and how these parameters are treated in the fit.

We perform fits to each data set where $2\beta_s$, R_S , δ_S , $R_{||}$, $\delta_{||}$, $R_{\perp}$, $\delta_{\perp}$ are free parameters and all other parameters are kept fixed. We also perform fits where the S-wave component is present in the generated toy data, but ignored in the fit (R_S is set to 0) in order to investigate the bias in the determination of $2\beta_s$.

The results of these fits for the statistical error and mean value of the weak phase $-2\beta_s$ are summarized in Table 3, 4 and 5 for several different scenarios with $-2\beta_s = -0.0368$, $-2\beta_s = -0.2$ and $-2\beta_s = -0.5$, respectively. As an example, in Figure 1 we show the distributions of the fitted values of $-2\beta_s$ for $R_S = 0.1$, $\delta_S = \pi/2$ and $-2\beta_s = -0.5$ for both the S-wave fraction R_S fixed to zero and R_S left free in the fits. In Figure 2 we show the distributions of the fitted values of R_S and the strong phase of the S-wave component δ_S for the same case with R_S left free in the fit. It can be seen that when all parameters are fitted the results are unbiased, but when it is wrongly assumed that $R_S = 0$, the result for $-2\beta_s$ acquires a bias with regard to the true input value.

Figure 3 shows the bias in $-2\beta_s$ from neglecting an S-wave component with $R_S = 0.1$ and $\delta_S = \pi/2$ versus the value of $-2\beta_s$ used to generate the data sets. A linear dependence is observed, which demonstrates that the bias in $-2\beta_s$ is proportional to the true value of $-2\beta_s$. From Tables 3, 4 and 5 we observe biases for these scenarios which range from 7 – 17% in the measurement of $2\beta_s$ if an S-wave component is present, but left unaccounted for in the fits. The bias moves the measured value of $2\beta_s$ towards zero. This implies that the neglected CP-odd S-wave contribution has a bigger probability to be mis-identified as the CP-even longitudinal or parallel components than as the CP-odd perpendicular component. Therefore, although the bias from neglecting an S-wave contribution is unlikely to lead to false signal of new physics, it will cause a loss of

sensitivity to new physics. On the other hand, including the S-wave in the fit removes the bias in the central value of $2\beta_s$ at a cost of an increase of less than 15% in the statistical error.

	Float R_S in fit	Fix R_S to 0 in fit
$R_S = 0$		$\sigma(2\beta_s) = 0.045$, $\text{Mean}(2\beta_s) = 0.038$
$R_S = 0.1, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.035$	$\sigma(2\beta_s) = 0.045$, $\text{Mean}(2\beta_s) = 0.032$
$R_S = 0.1, \delta_S = 0$	$\sigma(2\beta_s) = 0.054$, $\text{Mean}(2\beta_s) = 0.040$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.036$
$R_S = 0.05, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.040$	$\sigma(2\beta_s) = 0.045$, $\text{Mean}(2\beta_s) = 0.036$
$R_S = 0.05, \delta_S = 0$	$\sigma(2\beta_s) = 0.055$, $\text{Mean}(2\beta_s) = 0.038$	$\sigma(2\beta_s) = 0.047$, $\text{Mean}(2\beta_s) = 0.032$

Table 3: Statistical errors and mean values of $2\beta_s$ from 500 fits for different scenarios with $2\beta_s = 0.0368$. The errors on $\sigma(2\beta_s)$ and $\text{mean}(2\beta_s)$ are approximately 0.003 and 0.002, respectively. The same data sets are used to obtain the results in the second and third columns.

	Float R_S in fit	Fix R_S to 0 in fit
$R_S = 0$		$\sigma(2\beta_s) = 0.044$, $\text{Mean}(2\beta_s) = 0.198$
$R_S = 0.1, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.052$, $\text{Mean}(2\beta_s) = 0.199$	$\sigma(2\beta_s) = 0.047$, $\text{Mean}(2\beta_s) = 0.166$
$R_S = 0.1, \delta_S = 0$	$\sigma(2\beta_s) = 0.056$, $\text{Mean}(2\beta_s) = 0.202$	$\sigma(2\beta_s) = 0.049$, $\text{Mean}(2\beta_s) = 0.170$
$R_S = 0.05, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.049$, $\text{Mean}(2\beta_s) = 0.197$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.182$
$R_S = 0.05, \delta_S = 0$	$\sigma(2\beta_s) = 0.053$, $\text{Mean}(2\beta_s) = 0.198$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.180$

Table 4: Statistical errors and mean values of $2\beta_s$ from 500 fits for different scenarios with $2\beta_s = 0.2$. The errors on $\sigma(2\beta_s)$ and $\text{mean}(2\beta_s)$ are approximately 0.003 and 0.002, respectively. The same data sets are used to obtain the results in the second and third columns.

	Float R_S in fit	Fix R_S to 0 in fit
$R_S = 0$		$\sigma(2\beta_s) = 0.051$, $\text{Mean}(2\beta_s) = 0.501$
$R_S = 0.1, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.059$, $\text{Mean}(2\beta_s) = 0.501$	$\sigma(2\beta_s) = 0.053$, $\text{Mean}(2\beta_s) = 0.415$
$R_S = 0.1, \delta_S = 0$	$\sigma(2\beta_s) = 0.061$, $\text{Mean}(2\beta_s) = 0.501$	$\sigma(2\beta_s) = 0.052$, $\text{Mean}(2\beta_s) = 0.417$
$R_S = 0.05, \delta_S = \pi/2$	$\sigma(2\beta_s) = 0.051$, $\text{Mean}(2\beta_s) = 0.506$	$\sigma(2\beta_s) = 0.048$, $\text{Mean}(2\beta_s) = 0.463$
$R_S = 0.05, \delta_S = 0$	$\sigma(2\beta_s) = 0.053$, $\text{Mean}(2\beta_s) = 0.501$	$\sigma(2\beta_s) = 0.049$, $\text{Mean}(2\beta_s) = 0.461$

Table 5: Statistical errors and mean values of $2\beta_s$ from 500 fits for different scenarios with $2\beta_s = 0.5$. The errors on $\sigma(2\beta_s)$ and $\text{mean}(2\beta_s)$ are approximately 0.003 and 0.002, respectively. The same data sets are used to obtain the results in the second and third columns.




Figure 1: Distributions of the fitted values of $-2\beta_s$ for the scenario $R_S = 0.1, \delta_S = \pi/2, 2\beta_s = 0.5$. The left and right plots are obtained with or without fixing R_S to 0 in fitting the data, respectively.




Figure 2: Distributions of the fitted values of R_S and δ_S for the scenario $R_S = 0.1, \delta_S = \pi/2, 2\beta_s = 0.5$ without fixing R_S to 0 in fitting the data.




Figure 3: The bias in $-2\beta_s$ from neglecting an S-wave component with $R_S = 0.1$ and $\delta_S = \pi/2$ versus the value of $-2\beta_s$ used to generate the data sets. The bias is the difference of the mean of the fitted to the generated $-2\beta_s$ values. A linear fit is superimposed on the graph.

4 Measuring $\cos 2\beta_s$

In Eq. 2.1 and 2.2 one observes that the differential decay rates are invariant under the transformation

$$(\delta_{\parallel} - \delta_0, \delta_{\perp} - \delta_0, \delta_S - \delta_0, -2\beta_s, \Delta\Gamma_s) \leftrightarrow (\delta_0 - \delta_{\parallel}, \pi + \delta_0 - \delta_{\perp}, \delta_0 - \delta_S, \pi - (-2\beta_s), -\Delta\Gamma_s) . \quad (4.1)$$

As a consequence the measurement of $2\beta_s$ is subject to a two-fold ambiguity, which is equivalent to $\cos 2\beta_s$ transforming into $-\cos 2\beta_s$. A measurement of $\cos 2\beta_s$ including its sign would allow us to resolve this ambiguity.

If the interference between the P-wave and S-wave amplitudes were to be significant in the $\phi(1020)$ mass region, we could use this effect to measure $\cos 2\beta_s$, in the same way as BaBar measured $\cos 2\beta$ in $B^0 \rightarrow J/\psi K_S^0 \pi^0$ [13]. This requires measuring $\delta_S - \delta_0$, the strong phase difference between the S-wave and the longitudinal P-wave, as a function of the K^+K^- mass in the $\phi(1020)$ mass region. When plotting this function, two branches are expected with each corresponding to a different solution for the weak phase (see Figure 4 left). It is straightforward to choose the physical solution since the phase of the P-wave Breit-Wigner amplitude is expected to rise rapidly through the $\phi(1020)$ mass region (dashed red curve in Figure 4 right), while the phase of the S-wave amplitude, which can be described either by a coupled channel Breit-Wigner function in case of an f_0 contribution or by a constant term in case of a non-resonant contribution, is expected to vary relatively slowly (dotted green curve in Figure 4 right), resulting in $\delta_S - \delta_0$ rapidly falling with increasing K^+K^- mass (solid blue curves in Figure 4).




Figure 4: An example to illustrate the dependence of the strong phase of the S-wave δ_S , of the strong phase of the longitudinal P-wave δ_0 , and of their difference $\delta_S - \delta_0$, on the K^+K^- mass. Left: the solid blue curve is the physical solution for $\delta_S - \delta_0$ and the dashed black curve shows the mirror solution. Right: the dashed red, dotted green and solid blue curves are for δ_0 , δ_S , and $\delta_S - \delta_0$, respectively.

Below we use a Monte Carlo simulated toy data set to demonstrate the feasibility of this method in measuring the sign of $\cos 2\beta_s$. We generate 30000 $B_s^0 \rightarrow J/\psi K^+K^-$ events in the K^+K^- mass region between 1 and 1.05 GeV/c^2 , roughly corresponding to 0.5 fb^{-1} of integrated luminosity. The P-wave and f_0 contributions are included coherently. The values of the parameters used to generate the toy data set are the same as in Table 2 except that we set $-2\beta_s = -0.0368$, and that the values of both R_S and δ_S depend on

the K^+K^- mass. The f_0 contribution accounts for about 10% of the total decay rate in the given mass region, as is shown in Figure 5.




Figure 5: The data points correspond to the K^+K^- mass distribution of a generated sample of $B_s^0 \rightarrow J/\psi K^+K^-$ events including 10% f_0 contribution in the mass region. The dotted red curve indicates the f_0 contribution.

The data sample is divided into bins in the K^+K^- mass. For each bin i , two parameters $\delta_{S,i}$ and $R_{S,i}$ are used to represent the average strong phase and the fraction of the f_0 contribution. Both $\sin 2\beta_s$ and $\cos 2\beta_s$ are treated as independent free parameters. Common free parameters $\sin 2\beta_s$, $\cos 2\beta_s$, $R_{||}$, $R_{\perp}$, $\delta_{||}$, $\delta_{\perp}$, Γ_s and $\Delta\Gamma_s$ are used for all bins. Note that we still adopt the convention $\delta_0 = 0$ as only the relative phase differences in each bin can be measured. A combined fit to the time-dependent angular distributions of all the bins is performed to extract these free parameters. The fitted values of the strong phase difference $\delta_S - \delta_0$ versus the K^+K^- mass are plotted in Figure 6. The two branches correspond to opposite values of $\cos 2\beta_s$. Just as expected, the branch corresponding to the true solution decreases rapidly around the nominal $\phi(1020)$ mass. Choosing this branch leads to the unique solution

$$\sin 2\beta_s = 0.043 \pm 0.05, \quad \cos 2\beta_s = 1.05 \pm 0.08, \quad (4.2)$$

which gives the ambiguity-free result

$$-2\beta_s = -0.043 \pm 0.05. \quad (4.3)$$

In this example, the measured $-2\beta_s$ is separated from $\pi - (-2\beta_s)$ by 13σ , therefore the discrete ambiguity in $2\beta_s$ is completely resolved. Although the actual measurement precision in $\cos 2\beta_s$ will depend on the size of the f_0 contribution as well as background, the possibility to resolve the ambiguity in $-2\beta_s$ using this method is very promising.

5 Conclusions

In the decay $B_s^0 \rightarrow J/\psi K^+K^-$ we expect that a K^+K^- S-wave contribution in the narrow $\phi(1020)$ mass region could be as large as 10%. The full differential decay rates for this decay including the S-wave contribution have been presented. We have considered a range




Figure 6: The fitted values of $\delta_S - \delta_0$ versus K^+K^- mass are shown in red and black data points, corresponding to opposite values of $\cos 2\beta_s$. The blue curve shows the dependence of $\delta_S - \delta_0$ on K^+K^- mass implemented in simulation.

of scenarios which include S-wave components of 5% and 10%. We have shown that within these scenarios, if an S-wave component is ignored in the analysis, the measurement of the weak phase $-2\beta_s$ would be biased by between 7% and 17% towards zero. We have demonstrated that by properly allowing for this S-wave component in the fit, an unbiased measurement of $2\beta_s$ may be obtained with a slightly increased statistical error. Finally, we have shown that the interference between the K^+K^- S-wave and P-wave amplitudes can be used to resolve the two-fold ambiguity in the measurement of the weak phase $-2\beta_s$.

Acknowledgments

The authors would like to acknowledge the LHCb colleagues for useful and stimulating discussions. We particularly thank Tim Gershon, Olivier Leroy and Guy Wilkinson for valuable suggestions.

References

- [1] M. Artuso et al., *B, D and K decays*, Eur. Phys. J. **C 57** (2008) 309 [arXiv:0801.1833].
- [2] P. Ball and R. Fleischer, *Probing new physics through B mixing: status, benchmarks and prospects*, Eur. Phys. J. **C 48** (2006) 413 [hep-ph/0604249].
- [3] P. Clarke, C. Mclean and A. Osorio-Oliveros, *Sensitivity studies to β_s and $\Delta\Gamma_s$ using the full $B_s^0 \rightarrow J/\psi\phi$ angular analysis at the LHCb*, CERN-LHCB-2007-101.
- [4] M. Ciuchini et al., *Next-to-leading order strong interaction corrections to the $\Delta(F) = 2$ effective hamiltonian in the MSSM*, JHEP **09** (2006) 013 [hep-ph/0606197].
- [5] A. Lenz and U. Nierste, *Theoretical update of $B_s - \bar{B}_s$ mixing*, JHEP **06** (2007) 072 [hep-ph/0612167].
- [6] Z. Ligeti, M. Papucci and G. Perez, *Implications of the measurement of the $B_s^0 - \bar{B}_s^0$ mass difference*, Phys. Rev. Lett. **97** (2006) 101801 [hep-ph/0604112].
- [7] CDF collaboration, T. Aaltonen et al., *First flavor-tagged determination of bounds on mixing-induced CP violation in $B_s^0 \rightarrow J/\psi\phi$ decays*, Phys. Rev. Lett. **100** (2008) 161802.
- [8] D0 collaboration, V. M. Abazov et al., *Measurement of B_s mixing parameters from the flavor-tagged decay $B_s^0 \rightarrow J/\psi\phi$* , Phys. Rev. Lett **101** (2008) 241801 [arXiv:0802.2255].
- [9] BABAR collaboration, B. Aubert et al., *Measurement of CP-violating asymmetries in the $B^0 \rightarrow K^+K^-K_S^0$ dalitz plot* [arXiv:0808.0700].
- [10] BABAR collaboration, B. Aubert et al., *Amplitude analysis of the decay $D^0 \rightarrow K^-K^+\pi^0$* , Phys. Rev. **D 76** (2007) 011102 [arXiv:0704.3593].
- [11] S. Stone and L. Zhang, *S-waves and the Measurement of CP Violating Phases in B_s Decays*, Phys. Rev. **D 79** (2009) 074024 [arXiv:0812.2832].
- [12] M. Calvi, O. Leroy and M. Musy, *Flavour tagging algorithms and performances in LHCb*, CERN-LHCB-2007-058.
- [13] BABAR collaboration, B. Aubert et al., *Ambiguity-free measurement of $\cos 2\beta$: Time-integrated and time-dependent angular analyses of $B \rightarrow J/\psi K\pi$* , Phys. Rev. **D 71** (2005) 032005 [hep-ex/0411016].